

# ExCell Bio

## OptiViro<sup>®</sup> NK Cell Expansion Basic Kit P01

For Research and Manufacturing Use

Not Intended for Diagnostic and Therapeutic Use

### User Manual

Catalog Number	NE000-N032
Catalog Number	NE000-N031
Catalog Number	NE000-N031S



---

## | PRODUCT DESCRIPTION

OptiVibro® NK Cell Expansion Basic Kit P01 is a serum-free, and xeno-free culture system specifically designed for the robust expansion of activated natural killer (NK) cells. This comprehensive kit includes:

- **OptiVibro® NK Cell Serum-free Basal Medium P01:** Providing a foundational medium for NK cell growth.
- **OptiVibro® Immune Cell Serum-free Medium Supplement UE01:** Enhancing the basal medium with essential nutrients.
- **OptiVibro® Cytokine III:** A crucial component for supporting NK cell proliferation.

This kit is meticulously formulated to support the selective expansion of NK cells derived from human peripheral blood mononuclear cells (PBMC) and umbilical cord blood mononuclear cells. It is also compatible with NK cells differentiated from induced pluripotent stem cells (iPSC) and established NK cell lines, making it a versatile solution for various research and manufacturing applications.

### **Key Features:**

- **Activation Compatibility:** The kit is designed to work seamlessly with OptiVibro® Cytokine I and Cytokine II for the activation of NK cells, ensuring a streamlined process from activation to expansion.
- **Alternative Activation Methods:** While formulated to work with our cytokines, the kit is also adaptable for use with alternative methods of NK cell activation, providing flexibility for various experimental designs.
- **Expansion of Activated NK Cells:** Primarily supports the expansion of already activated NK cells, making it an ideal choice for applications where NK cells have been pre-activated through different stimulation methods.
- **iPSC-derived NK Cells:** Effective for the expansion of NK cells derived from iPSC differentiation, facilitating research into immune cell therapy and regenerative medicine.
- **NK Cell Lines:** Suitable for the expansion of established NK cell lines, like NK92, supporting consistent and reliable results in immunological research and bioproduction.

## | SPECIFICATION, STORAGE AND TRANSPORTATION

### **REQUIREMENT**

Product Name	Cat.#	Specification	Storage	Transportation	Shelf Life
<b>OptiVibro® NK Cell Expansion Basic Kit P01</b>	<b>NE000-N032</b>	<b>1000 mL kit</b>	<b>2-8°C Protect From Light</b>	-	-
OptiVibro® NK Cell Serum-free Basal Medium P01	BA0092	1000 mL	2-8°C Protect From Light	< 25°C Protect From Light	12 months
OptiVibro® Immune Cell Serum-free Medium Supplement UE01	BA0332	8 mL	2-8°C Protect From Light	< 25°C Protect From Light	18 months
OptiVibro® Cytokine III	BA0132	310 µL	-20°C	< 0°C Protect From Light	12 months
<b>OptiVibro® NK Cell Expansion Basic Kit P01</b>	<b>NE000-N031</b>	<b>500 mL kit</b>	<b>2-8°C Protect From Light</b>	-	-
OptiVibro® NK Cell Serum-free Basal Medium P01	BA0091	500 mL	2-8°C Protect From Light	< 25°C Protect From Light	12 months
OptiVibro® Immune Cell Serum-free Medium Supplement UE01	BA0331	4 mL	2-8°C Protect From Light	< 25°C Protect From Light	18 months
OptiVibro® Cytokine III	BA0131	155 uL	-20°C	< 0°C Protect From Light	12 months
<b>OptiVibro® NK Cell Expansion Basic Kit P01</b>	<b>NE000-N031S</b>	<b>100 mL kit</b>	<b>2-8°C Protect From Light</b>	-	-
OptiVibro® NK Cell Serum-free Basal Medium P01	BA0091S	100 mL	2-8°C Protect From Light	< 25°C Protect From Light	12 months
OptiVibro® Immune Cell Serum-free Medium Supplement UE01	BA0331S	0.8 mL	2-8°C Protect From Light	< 25°C Protect From Light	18 months
OptiVibro® Cytokine III	BA0131S	31 uL	-20°C	< 0°C Protect From Light	12 months

---

## | PERFORMANCE, APPLICATION AND HANDLING

### RECOMMENDATIONS

1. Store cell culture medium in a dark environment, preferably in colored packaging, to protect from light exposure.
2. Avoid prolonged exposure to lighting during transport to prevent discoloration.
3. Implement thorough cleaning and sterilization methods for transport to sterile areas; avoid UV sterilization.
4. Switch off UV lamps when transferring through UV-sterilized windows.

## | EXPERIMENTAL MATERIALS AND REAGENTS

1. Peripheral blood mononuclear cells (PBMC) or umbilical cord blood mononuclear cells.
2. Heat-inactivated autologous plasma (commercial serum substitutes or human AB serum can also be used).
3. Culture plates/culture bottles/culture bags.
4. Lymphocyte separation fluid, DPBS solution or saline, centrifuge tubes, pipettes, pipette guns, and pipette tips.
5. CO<sub>2</sub> incubator, centrifuge, cell counter, inverted microscope, water bath, etc.

## | INSTRUCTION FOR USE

### **Prepare media**

1. Equilibrate OptiVibro® NK Cell Serum-free Basal Medium P01 and OptiVibro® Immune Cell Serum-Free Medium Supplement UE01 to room temperature for 1-4 hours.
2. In a biosafety cabinet, add 8 mL/4 mL of supplement to every 1 L/500 mL of basal medium. Mix by inverting 3-5 times to obtain the complete medium.
3. Add one vial of 310 µL/155 µL Cytokine III to every 1000 mL/500 mL of complete medium to prepare the NK cell expansion medium. The medium is stable for 3 weeks after preparation.

### **Activation and expansion of NK cells from PBMC**

## 1. Day0

**Flask Preparation:** If necessary for your specific NK cell activation protocol, coat the T75 culture flask as required.

**Cell Seeding:** In the T75 flask, combine NK culture medium, 10% heat-inactivated autologous plasma (1.5 mL), and any additional NK activation reagents as needed. Seed the PBMCs into the flask to achieve a total volume of 15 mL. Gently shake the flask to distribute the cells evenly and place it in a 37°C, 5% CO<sub>2</sub> incubator.

### 【Note】

Recommended starting cell density for PBMC seeding is 2-2.5×10<sup>6</sup> cells/mL. For cord blood with low initial NK ratio, increase to 3×10<sup>6</sup> cells/mL. Use an electric pipette to seed cells, avoiding contact with the coating and spreading evenly.

## 2. Day3

Slowly add 13.5 mL of NK culture medium and 10% heat-inactivated autologous plasma (1.5 mL) along the side wall of the culture flask.

## 3. Day5

Sample and count, adjust cell density to 1.0×10<sup>6</sup> cells/mL, and transfer to a T175 culture flask.

## 4. Day 7 and beyond

Sample and count every 1-2 days, adjust cell density to 0.5-1.0×10<sup>6</sup> cells/mL, and expand or transfer to a cell culture bag as needed. Reduce heat-inactivated autologous plasma to 1% from Day 7.

## 5. Harvest cells

Harvest cells on days 14-18.

## | OTHERS

1. If NK cells are isolated from PBMCs before culturing, the seeding density can be 1.0-2.0×10<sup>6</sup> cells/mL.

2. Inoculation density lower than 1.0×10<sup>6</sup> cells/mL may lead to culture failure.

## | DISCLAIMER

1. Use the product according to the manual instructions. Deviations from these instructions are at the user's risk.
2. This product is for scientific research and commercial production only and is not intended for clinical diagnosis or treatment. Users assume all risks for unauthorized use.